

What is being claimed is:

1. A method of expressing PCA interacting partners in plant material comprising:

(A) transforming said material with:

(1) a first construct coding for a first fusion product comprising

(a) a first fragment of a first molecule whose fragments can exhibit a detectable activity when associated and

(b) a first protein-protein interacting domain; and

(2) a second construct coding for a second fusion product comprising

(a) a second fragment of said first molecule and

(b) a second protein-protein interacting domain that can bind (1)(b);

(B) culturing said material under conditions allowing expression of said PCA interacting partners, and

(C) detecting said activity.

2. The method of claim 1 wherein the plant material is selected from the group consisting of whole plants and plant-derived organs, tissues, cells, subcellular parts, and protoplasts.

3. The method of claim 1 or claim 2 wherein the plant material is derived from a transgenic plant.

4. The method of claim 1 where an inducer is added to facilitate the interaction of said protein-protein interaction domains.

5. The method of claim 1 or 4 wherein a fluorescent substrate is added and said activity is detected using fluorescence microscopy, spectrofluorometry, FACS analysis, or a fluorescence-detecting video system.

5 6. The method of claim 1 or 4 where said plant material is cultured on a selective medium.

7. A system for use as a standard or control in a PCA assay or for use in validating a PCA assay comprising:

(a) a first fusion product comprising a fragment of a first molecule whose fragments can exhibit a detectable activity when associated and a first protein-protein interacting domain; and

(b) a second fusion product comprising a second fragment of said first molecule and a second protein-protein interaction domain that interacts with said first protein-protein interaction domain.

8. A system according to claim 7 where said first and second protein-protein interaction domains are selected from the group consisting of:

- 1) NPR1 + TGA2,
- 2) FKBP + FRB,
- 3) leucine zippers.

9. A PCA assay using any of the systems of claims 7 or 8.

10. A plant-based PCA assay using any of the systems of claims 7 or 8.

12. A plant transgenic for one or more genes, each independently selected from the group consisting of:

(A) 1 or more genes coding for 1 or more interacting partners able to participate in a PCA assay, and

5 (B) 1 more more genes which result, either directly or indirectly, in the presence of 1 or more interacting partners able to participate in a PCA assay.

12. A plant according to claim 12 where the plant is of the genus *Arabidopsis*.

10 13. A plant according to claim 13 where the plant is *Arabidopsis thaliana*.

14. A plant according to claim 12 where the interacting partners comprise one or more of a leucine zipper/reporter molecule fusion, a NPR1/reporter molecule fusion, a TGA2/reporter molecule fusion, a FKBP/reporter molecule fusion or, a FRB/reporter molecule fusion.

15 15. A method of determining whether a mutated gene acts upstream in a pathway affecting an inducible interaction comprising performing a PCA assay in a mutated plant and correlating a change in PCA activity, relative to that measured in a non-mutated control plant, with the presence of one or more genes acting upstream in said pathway.

20 16. A method of identifying one or more genes involved in a pathway controlling an inducible interaction which results in a monitorable activity comprising:

(1) mutagenizing a seed from a transgenic plant expressing an interacting partner involved in PCA;

(2) germinating the seed;

(3) treating with an inducer that controls the interaction of any interacting partners

present, and

(4) monitoring said activity, and

5 (5) correlating said activity with 1 or more genes involved in a pathway controlling an inducible interaction .

17

18. A method of cloning a gene comprising:

1) identifying a gene according to the method of claim 17, and

10 2) cloning said gene.

18

19. A gene cloned by the method of claim 18 or a gene substantially similar thereto.

19

20. A product derived from any of the genes of claim 19.

20

21. A vector comprising the gene of claim 19 or its product.

21

22. Biological material genetically transformed with the gene of claim 19 or the vector of claim 21.

22

20 23. A method comprising mutating a plant or plant material that exhibits a first level of interaction between PCA interacting partners and selecting for a resultant plant or plant material that exhibits a lower level of said interaction.

23
24. A plant or plant material formed according to the method of claim 23.

24
25. A method of identifying plant molecule that functions as a PCA interacting partner in a PCA assay comprising

5 (1) reacting

(A) a library of plant molecules which are fused to a first fragment of a reporter molecule, said first fragment exhibiting low or no activity, with

(B) a bait molecule fused to a second fragment of said reporter molecule, said second fragment also exhibiting low or no activity and

10 (2) correlating reconstitution of reporter molecule activity with the presence of a PCA interacting partner.

25
26. A method employing a Protein Complementation assay/Universal Reporter System (PCA/URS) for detecting and screening for ligands and/or bioregulators of a plant cellular receptor, which method comprises:

a) generating a first nucleic acid vector encoding a first fusion product comprising:

i) a first fragment of a first PCA/URS reporter molecule, and

ii) a second molecule, fused to said first fragment, which comprises a first subdomain of a cellular receptor molecule of interest;

20 b) generating a second nucleic acid vector encoding a second fusion product comprising:

i) a second fragment of said first PCA/URS reporter molecule, and

ii) a third molecule, fused to said second fragment, which comprises a second subdomain of said cellular receptor, and where said second subdomain may be the same as said first subdomain in the case of a homodimeric cellular receptor, or different from said first

subdomain in the case of a heterodimeric cellular receptor; or a receptor coactivator or a protein;

c) transfecting eukaryotic cells with said first and second nucleic acid vectors; and

d) testing said transfected cells for activity of said PCA/URS reporter molecule, said activity indicating reassociation of the first and second fragments of the PCA/URS reporter molecule mediated by the interaction of said first and second subdomains of the cellular receptor molecule; said association being induced by binding said receptor to said ligand or bioregulator.

27. A method employing a Protein Complementation Assay/Universal Reporter System (PCA/URS) for detecting and screening for ligands and/or bioregulators of a plant cellular receptor, which method comprises:

a) generating a first nucleic acid vector encoding a first fusion product comprising:

i) a first fragment of a first PCA/URS reporter molecule, and

ii) a second molecule, fused to said first fragment, which comprises a first subdomain of a cellular receptor molecule of interest;

b) generating a second nucleic acid vector encoding a second fusion product comprising:

i) a second fragment of said first PCA/URS reporter molecule, and

ii) a third molecule, fused to said second fragment, which comprises a second subdomain of said cellular receptor, and where said second subdomain may be the same as said first subdomain in the case of a homodimeric cellular receptor, or different from said first subdomain in the case of a heterodimeric cellular receptor;

c) transfecting eukaryotic cells with said first and second nucleic acid vectors;

d) obtaining a clonal population of cells that express said first and second fusion products; and

e) testing said transfected cells for activity of said PCA/URS reporter molecule, said

activity indicating reassociation of the first and second fragments of the PCA/URS reporter molecule mediated by the interaction of said first and second subdomains of the cellular receptor molecule; said association being induced by binding said receptor to said ligand or bioregulator.